

Success rate of miniscrew and miniplates as anchorage for orthodontic treatment: A Scoping Review

Dr. Pankaj Wadhwa¹, Dr. Pradeep Raghav², Dr. Shalu Jain³, Dr. Kumar Amit⁴, Dr. Prashant Sharma⁵

¹Post graduate student Department of orthodontics Subharti Dental College Swami Vivekanand Subharti University Meerut, Uttar Pradesh (India)

²Professor and Head Department of orthodontics Subharti Dental College Swami Vivekanand Subharti University Meerut, Uttar Pradesh (India)

³Associate professor Department of orthodontics Subharti Dental College Swami Vivekanand Subharti University Meerut, Uttar Pradesh (India)

⁴Associate professor Department of orthodontics Subharti Dental College Swami Vivekanand Subharti University Meerut, Uttar Pradesh (India)

⁵Assistant professor Department of orthodontics Subharti Dental College Swami Vivekanand Subharti University Meerut, Uttar Pradesh (India)

¹pankajwadhwa59@gmail.com

²drpradeeppraghav@gmail.com

³shalu.j07@gmail.com

⁴doctoramit2010@gmail.com

⁵drprashantsharma94@gmail.com



This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Introduction:

Conventional anchorage methods generally require patient corporation which may result in undesirable contrary movement of teeth and can be restricted in dentitions that are compromised. Miniscrew and miniplate are specifically designed for enhancing orthodontic anchorage and are being increasingly implemented into orthodontics. Objectives: To conduct a scoping review for the evaluation of the success rate of miniscrew and miniplates as anchorage unit during orthodontic treatment.

Methods: Strategic and detailed search of the literature in various databases was commenced till October 2020 utilizing free text and MeSH terms, followed by PRISMA to classify different studies for extraction of the data.

Results and conclusions: It was concluded that granulation tissue formation and rate of Inflammation was higher in miniplate than miniscrew, however statistical differences were not found. Another factor observed was the effect of mean age on the success rate of miniscrew and miniplate between the age group of 10-20 years and 20-30 years. It was found that the success rate of both miniscrew and miniplate were similar in both the age groups but as the age advanced from 10-20 years to 20-30 years, the success rate of miniscrews decreased while it remained constant for miniplates. Based on the length, it was seen that in both primary and secondary insertions 8.0-mm miniscrews were significantly safer than the 6.0-mm miniscrews whereas, for diameter, a screw of 1mm or less was associated with mobility or failure of the miniscrews. In relation to the site palatal implants in fixed appliance therapy. However, of placement in the posterior region miniscrews in the different factors like insufficient space, problems in maxilla was a higher success rate than that in the mandible. **Keywords:** Miniscrews, Miniplates, Success rate, Maxilla, Mandible.

Main Points:

1. Granulation tissue formation and the rate of Inflammation was higher in miniplates than miniscrews.
2. The success rate of both miniscrews and miniplates were similar in both the age groups but as the age advanced from 10-20 years to 20-30 years.
3. The success rate of miniscrews decreased with age from 20- 30 years while it remained constant for miniplates.
4. Based on the length, 8.0-mm miniscrews were significantly anterior en-masse retraction, molar uprighting, molar more stable than the 6.0-mm miniscrews in both primary and secondary insertions
5. Based on diameter, a screw of 1mm or less was associated with mobility or failure of the miniscrews.

I. Introduction

Anchorage is one of the demanding aspects of Orthodontics. Headgear, elastics, and a number of other appliances have been suggested as effective forms of Miniscrews can also be repositioned during treatment to allow all intended tooth movements to be accomplished orthodontic anchorage for since long. The use of Conventional anchorage methods generally requires patient

cooperation and often results in undesirable tooth enough to be placed in any space in the alveolar bone, even movements. Also, their use can be restricted in patients in interdental areas. However, there are failures as well like with compromised dentitions.¹ The incorporation of the fracture at the time of placement, loosening under loading, skeletal anchorage system into orthodontic treatment for the last 25 years has been able to overcome some of these and impingement on roots either during placement or during tooth movement. shortcomings of conventional anchorages.

Absolute anchorage needs have paved the way for the development of miniscrews and it all started with the use of conventional dental implants, retromolar implants, and palatal implants in fixed appliance therapy. However, different factors like insufficient space, problems in connecting to orthodontic attachments, and cost factor became limiting and this led to the development of smaller devices that could be placed in various locations in the dental arch.¹ Miniscrews and miniplates were designed for such purpose and are being progressively adopted into orthodontics practice for absolute anchorage needs. The indications for treatment with miniscrew implants include, to name a few, molar protraction, intrusion of supra-erupted teeth, intrusion of posterior in anterior open bite, 4. Based on the length, 8.0-mm miniscrews were significantly anterior en-masse retraction, molar uprighting, molar distalization, traction of impacted canine, and attachment of protraction facemask.²

Recent scientific literature is replete with articles about the application of miniscrews in orthodontics and their small size and apparent ease of placement add to their merit. Miniscrews can also be repositioned during treatment to allow all intended tooth movements to be accomplished and they have added advantages of low cost, simple surgical placement, and ease of removal. They are small enough to be placed in any space in the alveolar bone, even in interdental areas. However, there are failures as well like fracture at the time of placement, loosening under loading, skeletal anchorage system into orthodontic treatment for the last 25 years has been able to overcome some of these and impingement on roots either during placement or during tooth movement.

To overcome the limitations of miniscrews, surgical miniplates with intraoral attachments have been used as an alternative to serve as temporary skeletal anchorage devices (TSAD). Miniplates are associated with less fracture rate and risk of root impingement as they are placed at a safer distance

from the roots & also allow the free movement of roots over the TSAD. The initial stability and survival rate (“Miniscrews”) AND (“Miniplates”)) AND of orthodontic mini-implants are highly dependent on the (“Orthodontics”)/ (“Success rate”) AND (Miniscrew)) amount of cortical bone at their insertion site. In areas with AND (Miniplates)) AND (“Orthodontics”). limited bone availability, miniplates are preferred to This search strategy was applied to the key databases provide effective skeletal anchorage.³ However, the need for surgery and high cost are their disadvantages

Orthodontic literature has a good number of studies that have evaluated the success of miniscrews based on various parameters which are Granulation formation rate (GFR), Inflammation rate,⁴ Mean age & Site of Placement⁵ but very few researches have been conducted on the stability of miniplates. However, as per the limited studies available, miniplates are also said to be an equally effective method of enhancing the anchorage but since the number of studies based on the stability of miniscrews exceeds that of miniplates a judicious comparison between the two is not available so the main aim of this scoping review was to evaluate the success rate of miniscrews and miniplates as anchorage unit for orthodontic treatment based on Granulation formation rate, Inflammation rate, Mean age, and Site of placement.

II. Search Strategy And Selection Criteria

A scoping review of the literature was conducted to evaluate success rate of miniscrew and miniplates as anchorage for orthodontic treatment. The PRISMA search strategy was planned to use the MeSH terms and Boolean terminology:

((“Success Rate”) And (“Miniscrews”)) And (“Orthodontics”)/ ((“Success Rate”) And (“Miniplates”)) And (“Orthodontics”)/ (“Success Rate”) Or (“Miniscrews”) And (“Miniplates”)) And (“Orthodontics”)/ ((“Success Rate”) And (Miniscrew)) And (Miniplates)) And (“Orthodontics”).

This search strategy was applied to the key databases PUBMED, SCOPUS, WEB OF SCIENCE. Furthermore, the reference lists of the included studies and previous reviews were thoroughly searched to identify any potential articles to be included in this review. We restricted searches to trials in human participants with the full text published in English. We also searched Case reports, Grey literature, and unpublished literature in ClinicalTrials.gov, and full research articles were included except abstract for scientific presentation and review papers. The included

articles were screened thoroughly, and the level of evidence was determined based on Oxford Centre for Evidence-based Medicine (OCEBM).

Inclusion Criteria:

a) Patients Who Were Treated With Orthodontic treatment With The Use Of Miniscrews And Miniplates.

- Healthy patients from both genders and from 10 to 30 years of age.
- Optimum Oral Hygiene before treatment with no signs of gingivitis and/or periodontitis.

b) No restrictions on patient's sex, city, country, ethnicity, and socioeconomic status

Exclusion Criteria:

- Patients with poor oral hygiene.
- Those who are not treated with miniscrews and miniplates were excluded.
- Studies done on Animals were also excluded.

III. Study Design

- Included those studies in which the investigators used miniscrews and miniplates as anchorage for intervention.
- No restrictions were implemented regarding the follow-up time or the number of participants.
- Studies on the human population published in the English language were included.
- Systematic reviews, Meta-analysis, Randomized Controlled Trials, Case-Control, Cohort studies, Case reports, full research articles, and review papers were included except for the commentaries.

IV. Information Sources

The search was conducted by using the following databases

- PubMed / Medline
- Scopus

- Web of Science

V. Literature Search and Screening

The PRISMA search strategy was planned to use the MeSH terms and Boolean terminology: ((“Success rate”) AND (“Miniscrews”)) AND (“Orthodontics”)/ ((“Success rate”) AND (“Miniplates”)) AND (“Orthodontics”)/ (“Success rate”) OR (“Miniscrews”) AND (“Miniplates”)) AND (“Orthodontics”)/ ((“Success rate”) AND (Miniscrew)) AND (Miniplates)) AND (“Orthodontics”).

This search strategy was applied to the key databases PUBMED, SCOPUS, WEB OF SCIENCE. Furthermore, the reference lists of the included studies and previous reviews were thoroughly searched to identify any potential articles to be included in this review. We restricted searches to trials in human participants with the full text published in English. We also searched Case reports, Grey literature, and unpublished literature in English language were included. ClinicalTrials.gov, and full research articles were except abstract for scientific presentation and review Controlled Trials, Case-Control, Cohort studies, Case reports, full research articles, and review papers. The included articles were screened thoroughly, and the level of evidence was determined based on Oxford Centre for Evidence-based Medicine (OCEBM). (Table.1)

VI. Results

A.1 Study Selection:

There were a total of 1027 articles identified through the search strategy and sources listed previously. After removal of duplication 941 articles were taken. All the titles were screened, out of which a total of 855 articles were rejected as they were not matching the search question. On the screening of abstracts, 27 articles were excluded as they were not related to success rate, animal study, or not in the English language. The remaining 59 articles were read for a complete evaluation of the text out of which 31 articles were excluded as they were not meeting the inclusion criteria. Finally, 28 articles were selected for inclusion in this review for qualitative and quantitative synthesis. (Figure 1)




Figure 1. PRISMA Search Algorithm

A.2 Study Characteristics:

All Included articles were all in English Language and published between 2000 to 2020. The articles included were on subjects treated with miniscrews and miniplates as anchorage for orthodontic treatment. There was a total of 28 prospective and retrospective studies taken in this scoping review. (Table 2-6)

VII. Discussion

The aim of this scoping review was to evaluate the success rate of miniplates and miniscrews, as skeletal anchorage. These are the most effective bone anchor system method which is used in cases having inadequate dental anchorage required for optimum orthodontic tooth movement. It is well known that articles of prospective and retrospective study both can provide clinical evidence and form the source of a good quality scoping review. Amongst them, there were few articles that compared different factors between miniscrews and miniplates, based on the choice of the clinician and many other factors in choosing the type of anchorage method to be used.

Recent methods such as miniscrew have proved to be an alternative anchorage system. Through insertion, they are easily placed and removed and can be packed immediately. However, stability is limited after loading with torsion. Miniplates provide various advantages such as high anchorage, less root injury, high success rates, and less risk of fracture.

The aim of this scoping review is to discuss the success rate of miniscrew and miniplates in orthodontics patients. The selected studies are of different level of evidence which was used to answer the clinical question posed in this scoping review. This study included a review of factors (Granulation formation rate, Inflammation rate, Mean age & Site of Placement) related to the

success rate of miniscrews and miniplates installed in the maxilla and mandible in different regions.

Among Selected articles there was only one manuscript which compared miniscrew and miniplates in terms of granulation tissue formation rate and inflammation rate factors was considered for miniscrew and miniplates. In this study 0.9% of subjects showed granulation tissue formation while using miniplates, whereas only 0.6% showed granulation tissue when palatal screws were used and none of the subjects showed granulation tissue formation in the group where self drilling screws were replaced. Similarly, the rate of Inflammation was 7.6% in the miniplate group which was higher than 1.3% in the self drilling miniscrew group and 2.5% in the palatal screw group.

To study the effect of mean age as a significant factor in assessing the success rate of miniscrews and miniplates, the present study was segregated amongst two mean age groups, ranging from 10 to 20 years for early adolescents and 20 to 30 years for adults.

According to various studies focusing on 10-20 year age group, the overall success rate of miniscrew ranged from 77%-100%.^{20,21,23} However when miniplates were used for a similar age group, studies have shown a lesser variation with a mean survival rate of 93.6%.^{16,29} Total of 19 articles for miniscrew and 7 articles for miniplates were selected for evaluating the effect of overall success rate on mean age between 20 to 30 years.

As for the success of miniscrew, Hoi- Jeong Lim et al¹² in their study found that the overall success rate of miniscrews was 83.6% whereas the mandible had a more success rate of 0.48 times than that of the maxilla, similar results were found by Watanabe et al¹⁹ where in a superlative results were seen when the screws are loaded significantly higher success rate was observed in the maxilla(90.6%) when compared to the mandible(70.69%). The possible reason for this has been explained by Kuroda et al¹⁰ who concluded that the proximity of the miniscrew to the root surface is a potential threat for the failure of miniscrew anchorage and this is more evident in the mandible and thus the miniscrew failure rate is more in the mandible than that of the maxilla.

The stability of miniscrew is also dependent on its length, this has been stated by Maino et al.¹⁷ In another study by Suzuk et al,²² the authors took length as a prime factor to check the stability of miniscrew and found a significantly lower success rate with 5-mm miniscrews in comparison to the 6-mm and 7-mm miniscrews, showing a direct groups, ranging from 10 to 20 years for early adolescents correlation between the length of miniscrew and its stability. In a similar study, Uesugi et al²⁵ found 8.0-mm miniscrews to be significantly more stable than the 6.0-mm miniscrews. Thus, it can be concluded that a screw with a greater length tends to be more stable than a screw with smaller length.

Another factor that affects the stability of miniscrew is its diameter. G.Janson et al²¹ in this study found that a screw with a diameter of 1.0 mm or less was associated with the mobility or failure of the miniscrew. However other authors such as Maino et al¹⁷ have stated that the screw diameter resulted in no significant effects and rather the quality of bone has a greater role to play in the stability of miniscrew where soft bone quality and hard bone quality are the two factors for miniscrew failure, while the superlative results were seen when the screws are loaded into the bone of medium quality i.e. 10-15 Ncm. This has been more elaborately studied by Ichinohe et al²⁷ who concluded that mini-screws could be stabilized when the cortical bone thickness was ≥ 1.5 mm. The success rate was significantly higher in groups in which insertion depths of ≥ 4.5 mm. These results showed that sufficient cortical bone thickness is important for the primary stability of mini-screws with less impact on the screw diameter.

Takashi Watanabe et al²⁶ conducted a study and concluded that the overall success rate of miniscrew was 85.80%. In their study, one of the main factors analyzed was Mean insertion torque where the values were 10.7 ± 1.9 N.cm in the failure group and 8.5 ± 2.1 N.cm in the success group showing that one of the main causes of failure of Miniscrew is increased torque during insertion. Makoto Suzuki²² also obtained similar results in his investigation in which the miniscrews when it is placed with insertion torque greater than 10 Ncm had a lower success rate. This can be a possible explanation for a higher rate of miniscrew from the crest is more compatible with a denser miniscrew failure in the posterior part of the mandible due to the compact nature of bone in that region which tends to increase the insertion torque and leads to poor stability of miniscrew.

One reason which can affect the stability of miniscrew as stated by Hoi- Jeong Lim et al¹⁵ is the clinician's experience. They examined more than 20 miniscrews were placed by more experienced clinicians. The results have shown that there is an approximately 3.6-fold higher success rate of initial stability compared with those placed by less experienced clinicians after adjusting for the insertion site. Thus the results suggested that the initial stability depends on both, the clinician's experience as well as the insertion site.

A similar factor of less clinical importance has been studied extensively by Nikolaos Topouzelis et al¹⁸, who stated that factors such as the number of miniscrews used per patient and correlated with the success rate of conducted by Chung Ho et al⁸ who also confirmed that a miniscrews. It was noticed that the success rate decreased by 67% whenever supplementary miniscrew was used in the patient's oral cavity, They also evaluated the soft tissue type of placement resulted in a lower success rate whenever miniscrews were placed in movable mucosa.

With respect to the miniscrews placed on the palatal aspect, Young Ho Kim et al¹³ observed that the total showing that one of the main causes of failure of number of miniscrew is placed

for mid palatal resulted in an overall success rate was 90.80%. However, Ramzi Haddad et al³⁰ concluded that the distance to the alveolar crest was strongly associated with the long-term stability and success of orthodontic mini implants. More apical placement of the can be a possible explanation for a higher rate of miniscrew from the crest is more compatible with a denser and thicker bucco-lingual/palatal bone level and this also enhances the long term stability of miniscrew. 7 articles were selected in which miniplates were used by different authors for mean subject age between 20 to 30 years.

Contrary to the literature available for miniscrews, very few researches have been conducted on the stability of miniplates. However, as per the limited studies available miniplates are also said to be an equally effective method of enhancing anchorage. Of various parameters that were used to evaluate the stability of miniscrews, Shouichi Miyawaki et al⁶ concluded that there was no significant association between the success rate and variables such as screw length, location of implantation, surgical approach, loading (immediate or delayed), age or gender and the overall success rate was found to be 96.40% of the studied samples. Similar results were seen in the study per patient and correlated with the success rate of conducted by Chung Ho et al⁸ who also confirmed that there was no significant differences in the risk factors for failure of miniplates with an overall success rate of 95.5%.

Although miniplates have proven to be very stable, there are a few demerits that have been reported in a study conducted by Marie A. Cornelis,¹ who observed that the most common problems were postsurgical swelling which usually lasts 5 days on average, and also observed that more than a third of the patients experienced having cheek irritation initially but it lessened over time and resulted in 82% of the patients determines that the surgical number of other limitations were also found such as small experience was better.

Raymond Lam et al²⁴ also found soft tissue inflammation as the most common complication, which was amenable to focused oral hygiene and antiseptic rinses and also 40% of cases experienced mild complications.

Site of Placement: The placement site of miniscrews in the mandible has shown a significant difference in adults. Cheol-Hyun Moon⁵ concluded that dislodgement of the Orthodontic miniscrew implant occurred in the first two months after insertion, when placed in the posterior region of the maxilla & mandible and more than 90% of the failures occurred within the first 4 months. In another study, Makoto Suzuki et al²² concluded that the success rate of the miniscrews in the maxilla (93.4%) was higher as compared to that in the mandible (70.3%) when miniscrew was placed b/w maxillary second pre-molars and first molars in both maxilla & mandible. Although Shouichi Miyawaki et al⁶ evaluated the placement of miniplates and concluded that a significant association could not be detected between the success rate and the site of implantation.

The limitation of this scoping review was that various Comparative studies which were ideally required were not available which could have helped us in assessing various factors such as bone density, infection, smoking, and gender. Along with this maximum number of studies included were of Level 4 according to the level of evidence as Randomized Controlled Trials and systematic reviews were not available in the literature, and since studies of different levels of evidence were taken for this review a number of other limitations were also found such as small sample size for males as compared to females, subjects who have dropped out of the study during treatment, along with this comparison was not done in some studies which might have led us to a certain degree of bias in the present study.

VIII. Summary And Conclusion

For the purpose of orthodontic anchorage, miniscrews, and miniplates are being progressively adopted into orthodontic practice. However, there are associated factors that cause the failure of miniscrews, like fracture during placement, loosening over time, underloading, and impingement on roots either during placement or tooth movement. On the other hand, miniplates, which are placed at a safe distance from the roots offer the advantages of reduced risk of root impingement and allow free movement of roots past the temporary skeletal anchorage device. Hence, this study was conducted to evaluate the success rate of miniscrews Miyawaki et al [6] evaluated the placement of miniplates and concluded that a significant association could not be found and miniplates during orthodontic treatment.

It was concluded from this scoping review that granulation detected between the success rate and the site of tissue formation and the rate of Inflammation was higher in miniplates than in miniscrews, however, statistical differences were not found. The limitation of this scoping review was that various differences were not found. Another factor that was observed was the effect of mean age on the success rate of miniscrews and miniplates between the 10-20 years age group and 20-30 years age group. It was found that the both primary and secondary insertions whereas for success rate of both miniscrews and miniplates were similar in both the age groups but as the age advanced from 10-20 years to 20-30 years, the success rate of miniscrews decreased while it remained constant for miniscrews in the maxilla was higher than that in the miniplates. Based on the length, 8.0-mm miniscrews were significantly more stable than the 6.0-mm miniscrews in group and 20-30 years age group. It was found that the both primary and secondary insertions whereas for success rate of both miniscrews and miniplates were similar in both the age groups but as the age advanced from 10-20 years to 20-30 years, the success rate of diameter, a screw of 1mm or less was associated with mobility or failure of the miniscrews. In relation to the site of placement in the posterior region success rate of the miniscrews decreased while it remained constant for miniscrews in the maxilla was higher than that in the miniplates. Based on the length, 8.0-mm miniscrews were mandible.

Table 2. Evaluation Of Rate Of Granulation Tissue Formation &Inflammation Rate On Success Rate Of Miniscrews & Miniplates

S.NO	FACTORS	RESULT	OVERALL SUCCESS RATE
1	Granulation tissue formation rate	For Miniscrew- 0% For Miniplate- 0.9%	94% ⁴
2	Inflammation Rate	For Miniscrew- 1.3% For Miniplate- 7.6%	94% ⁴

Table 3. Evaluation Of Mean Age (From 10 To 20 Years) On Success Rate Of Miniscrew And Miniplates

S.NO	MEAN AGE FROM 10 TO 20 YEARS	MINISCREWS	MINIPLATES
1	MEAN AGE- 11.4 years	-----	Overall success- 93.6% ²⁰
2	MEAN AGE - 14.7 years	-----	Overall success- 100% ¹⁸
3	MEAN AGE- 16.99 year	Overall success- 90% ²¹	-----
4	MEAN AGE - 18.9 years	Maxilla-100%, Mandible-77.8% ²⁰	-----
5	MEAN AGE- 19.26 years	Maxilla-93.53%, Mandible-78.31% ²³	-----

Table 4. Evaluation Of Success Rate In Mean Age (From 20 To 30 Year) For Miniscrews And Miniplates As Anchorage Unit.

S.NO	MEAN AGE FROM 20 TO 30	MINISCREWS OVERALL SUCCESS RATE	MINIPLATES OVERALL SUCCESS RATE
1	MEAN AGE- 20.9 years	Maxilla- 93.4%, mandible- 70.3% ²²	-----
2	MEAN AGE- 21 years	maxilla- 90.6%, mandible - 70.69% ¹⁹	-----
3	MEAN AGE- 21.4years	Overall success- 64% ¹¹	-----
4	MEAN AGE- 21.8 years	Overall success- 84.4 % ⁶	-----
5	MEAN AGE- 21.8 years	-----	Overall success- 96.40% ⁶
6	MEAN AGE- 21.8 years	-----	Overall success- 86.8% ⁹
7	MEAN AGE- 21.8years	Overall success- 84.85% ⁹	-----
8	MEAN AGE- 21.9years	maxilla-86%, mandible-76.1 % ¹²	-----
9	MEAN AGE- 22.5years	Maxilla-89.7%, Mandible- 68.3% ¹⁰	-----
10	MEAN AGE- 23 years	maxilla- 93.1%, mandible- 93.2% ¹⁵	Overall success- 92.5% ¹
11	MEAN AGE- 23.2years	Overall success- 81% ¹⁴	-----
12	MEAN AGE- 23.4 years	Male-71.2%, Female-72.2% ²⁷	Overall success- 94% ⁴
13	MEAN AGE- 23.4 years	Overall success- 88.1% ¹⁰	-----
14	MEAN AGE- 23.4years	Overall success- 90.8% ¹³	-----
15	MEAN AGE- 23.7 years	-----	Overall success- 95.5% ⁸
16	MEAN AGE- 24.6 years	Overall success- 91.4% ¹⁷	-----
17	MEAN AGE- 25.4 years	Overall success- 85.8% ²⁶	-----
18	MEAN AGE- 25.7 years	-----	-----
19	MEAN AGE- 25.7years	Overall success- 94% ⁴	-----
20	MEAN AGE- 27.2 years	Overall success- 90.2% ¹⁸	-----
21	MEAN AGE- 27.5 years	-----	-----
22	MEAN AGE- 27.9 years	Primary: MB-79.1% MP-84.5% ²⁸ Secondary: MB-58.1% MP- 88.9% ²⁸	----- Overall success- 89% ⁷ Overall success- 98.6% ²⁴
23	MEAN AGE- 28.1 years	Primary-80.4%, Secondary-44.2% ²⁵	-----
24	MEAN AGE- 29 years	Overall success- 89% ⁷	-----
25	MEAN AGE- 29 years	-----	-----
26	MEAN AGE- 29.4 years	-----	-----

Table 5. Comparison Of Success Rate According To Site Of Placement Between Miniscrews And Miniplates

S.NO	SITE OF PLACEMENT	MINISCREWS OVERALL SUCCESS RATE	MINIPLATES OVERALL SUCCESS RATE
1	Posterior region of maxilla & mandible	Maxilla- 83.5%, Mandible- 84.1% ⁵	-----
2	Maxilla & Mandible- B/W maxillary second pre-molars & first molars	Maxilla- 93.4%, Mandible- 70.3% ²²	-----
3	Posterior region of maxilla & mandible	-----	Posterior Region Upper - 84.1% Posterior Region Lower- 81.6% ⁶

Table.6 Results Of The Reviewed Articles

S.NO	FACTORS	RESULTS SIGNIFICANT TO THE RESEARCH QUESTION
1.	Granulation tissue formation rate and Inflammation rate	More in miniplates as compared to miniscrews ⁴
2.	Mean Age (from 10-20 years)	Overall success rate similar in miniscrews and miniplates ^{16,20,21,23,29}
3.	Mean age (from 20-30 years)	Overall success rate decreased in miniscrews ^{22,19,11,6,9,12,10,15,14,27,30,13,17,26,18,4,28,15,7} Similar overall success rate in miniplates ^{6,9,14,8,7,24} there is no statistical difference was found between miniscrews and miniplates.
4.	Site of placement	Posterior region overall success rate of the miniscrews in the maxilla was higher than that in the mandible ^{5,22}

Table 1: Oxford centre for Evidence-Based Medicine 2011 Levels of Evidence

Question	STEP 1 (LEVEL 1*)	STEP 2 (LEVEL 2*)	STEP 3 (LEVEL 3*)	STEP 4 (LEVEL 4*)	STEP 5 (LEVEL 5)
How common is the problem?	Local & current random sample surveys	A systematic review of surveys that allow matching to local circumstances**	Local non-random sample**	Case series**	n/a
Is this diagnostic or monitoring test accurate? (diagnosis)	A systematic review of cross-sectional studies with consistently applied references standard & blinding	Individual cross-sectional studies with consistently applied references standard & blinding	Non-consecutive studies without consistently applied references standard**	Case-control studies or poor or non-independent reference standard**	Mechanism-based reasoning
What will happen if we do not add a therapy (prognosis)	A systematic review of inception cohort studies	Inception cohort studies	Cohort studies or control arm of randomized trial*	Case series or case control studies, poor quality prognostic cohort studies**	n/a
Does this intervention help? (treatment benefits)	Systematic review randomized trials or n-of 1 trial	Randomized trial or observational study with dramatic effect	Non-randomized controlled cohort/follow up study**	Case series or case control studies, historically control studies**	Mechanism based reasoning
What are the common harms? (treatment harms)	Systematic review randomized trials, Systematic review of nested case control studies	Individual Randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow up study provided there are sufficient number to rule out common harm**	Case series or case control studies, historically control studies**	Mechanism based reasoning
What are the rare harms? (treatment harms)	Systematic review randomized trials or n-of 1 trial	Randomized trial or (exceptionally) observational study with dramatic effect	Non-randomized controlled cohort/follow up study provided there are sufficient number to	Case series or case control studies, historically control studies**	Mechanism based reasoning

Is this (early detection) test worthwhile? (Screening)	Systematic review randomized trials	Randomized trial	Non-randomized controlled cohort/follow up study**	Case series or case control studies, historically control studies**	Mechanism based reasoning
--	-------------------------------------	------------------	--	---	---------------------------

*If not marked down on the basis of study quality, imprecision, indirectness (study PICO does not match questions PICO)

*Level may be graded down on the basis of study quality, imprecision, indirectness (study PICO does not match questions PICO), because of inconsistency between studies, or because the absolute effect size is very small; Level may be graded up if there is a large or very large effect sizes ** As always, a systematic review is generally better than an individual study.

References

- [1] Cornelis MA, Scheffler NR, Nyssen-Behets C, De Clerck HJ, Tulloch JC. Patients' and orthodontists' perceptions of miniplates used for temporary skeletal anchorage: a prospective study. Am J Orthod Dentofacial Orthop. 2008;133(1):18-24.
- [2] Chang HP, Tseng YC. Miniscrew implant applications in contemporary orthodontics. Kaohsiung J Med Sci. 2014;30(3):111-
- [3] Hourfar J, Kanavakis G, Goellner P, Ludwig B. Fully customized o p. placement of orthodontic miniplates: A technique. Head Face Med. 2014;10(1):14-8.

- [4] Takaki T, Tamura N, Yamamoto M, Takano N, Shibahara T, Yasumura T et al. Clinical study of temporary anchorage devices for orthodontic treatment. Bull Tokyo Dent Coll. 2010;51(3):151-63.
- [5] Moon CH, Lee DG, Lee HS, Im JS, Baek SH. Factors associated with the success rate of orthodontic miniscrews placed in the upper and lower posterior buccal region. Angle Orthod. 2008;78(1):101-6.
- [6] Miyawaki S, Koyama I, Inoue M, Mishima K, Sugahara T, Takano Yamamoto T. Factors associated with the stability of titanium screws placed in the posterior region for orthodontic anchorage. Am J Orthod Dentofacial Orthop. 2003;124(3):373-8.
- [7] Cheng SJ, Tseng IY, Lee JJ, Kok SH. A prospective study of the risk factors associated with failure of mini-implants used for orthodontic anchorage. Int J oral Maxillofac Implants. 2004;19(2):100-06.
- [8] Chen CH, Hsieh CH, Tseng YC, Huang IY, Shen YS, Chen CM. The use of miniplat osteosynthesis for skeletal anchorage. Plast Reconstr Surg. 2007;120(1):232-5.
- [9] Kuroda S, Sugawara Y, Deguchi T, Kyung HM, Takano Yamamoto T. Clinical use of miniscrew implants as orthodontic anchorage: success rates and postoperative discomfort. Am J Orthod Chang HP, Tseng YC. Miniscrew implant applications in Dentofacial Orthop. 2007; 131(1):9-15
- [10] Kuroda S, Yamada K, Deguchi T, Hashimoto T, Kyung HM, Yamamoto TT. Root proximity is a major factor for screw failure in orthodontic anchorage. Am J Orthod Dentofacial Orthop. 2007;131(4):68-73.
- [11] Justens E, De Bruyn H. Clinical outcome of mini-screws used as orthodontic anchorage. Clin Implant Dent Relat Res. 2008;10(3):174-80.
- [12] Lim HJ, Eun CS, Cho JH, Lee KH, Hwang HS. Factors associated with initial stability of miniscrews for orthodontic treatment. Am J Orthod Dentofacial Orthop. 2009;136(2):236-42.
- [13] Kim YH, Yang SM, Kim S, Lee JY, Kim KE, Gianelly AA et al. Midpalatal miniscrews for orthodontic anchorage: factors affecting clinical success. Am J Orthod Dentofacial Orthop. 2010;137(2):66-72.

- [14] Manni A, Cozzani M, Tamborrino F, De Rinaldis S, Menini A. Factors influencing the stability of miniscrews. A retrospective study on 300 miniscrews. *Eur J Orthod.* 2011;33(4):388-95.
- [15] Lim HJ, Choi YJ, Evans CA, Hwang HS. Predictors of initial stability of orthodontic miniscrew implants. *Eur J Orthod.* 2017;152(2):224-31. 2011;33(5):528-32.
- [16] Enache AM, Nicolescu I, Georgescu CE. Mandibular second molar impaction treatment using skeletal anchorage. *Rom J Morphol Embryol.* 2012;53(4):1107-10.
- [17] Maino BG, Pagin P, Di Blasio A. Success of miniscrews used as anchorage for orthodontic treatment: analysis of different factors. *Prog Orthod.* 2012;13(3):202-9.
- [18] Topouzelis N, Tsaousoglou P. Clinical factors correlated with the success rate of miniscrews in orthodontic treatment. *Eur J Oral Sci.* 2012;4(1):38-44.
- [19] Watanabe H, Deguchi T, Hasegawa M, Ito M, Kim S, Takano-Yamamoto T. Orthodontic miniscrew failure rate and root proximity, insertion angle, bone contact length, and bone density. *Orthod Craniofac Res.* 2013;16(1):44-55.
- [20] Samrit V, Kharbanda OP, Duggal R, Seith A, Malhotra V. Bone density and miniscrew stability in orthodontic patients. *Aust Orthod J.* 2012;28(2):204-12.
- [21] Janson G, Gigliotti MP, Estelita S, Chiqueto K. Influence of miniscrew dental root proximity on its degree of late stability. *Int. J. Oral Maxillofac.* 2013;42(4):527-34.
- [22] Suzuki M, Deguchi T, Watanabe H, Seiryu M, Iikubo M, Sasano T et al. Evaluation of optimal length and insertion torque for miniscrews. *Am J Orthod Dentofacial Orthop.* 2013;144(1):251-9.
- [23] Jing Z, Wu Y, Jiang W, Zhao L, Jing D, Zhang N et al. Factors Affecting the Clinical Success Rate of Miniscrew Implants for Orthodontic Treatment. *Int J Oral Maxillofac Implants.* 2016;31(4):12-5.
- [24] Lam R, Goonewardene MS, Allan BP, Sugawara J. Success rates of a skeletal anchorage system in orthodontics: A retrospective analysis. *Angle Orthod.* 2018;88(1):27-34.
- [25] Uesugi S, Kokai S, Kanno Z, Ono T. Prognosis of primary and secondary insertions of orthodontic miniscrews: what we have learned from 500 implants. *Am J Orthod Dentofacial Orthop* 2017;152(2):224-31.

- [26] Watanabe T, Miyazawa K, Fujiwara T, Kawaguchi M, Tabuchi M, Goto S. Insertion torque and Periotest values are important factors predicting outcome after orthodontic miniscrew placement. *Am J Orthod Dentofacial Orthop*. 2017;152(1):483-8.
- [27] Ichinohe M, Motoyoshi M, Inaba M, Uchida Y, Kaneko M, Matsuike R et al. Risk factors for failure of orthodontic mini-screws placed in the median palate. *J Oral Sci*. 2019;61(1):13-8.
- [28] Uesugi S, Kokai S, Kanno Z, Ono T. Stability of secondarily inserted orthodontic miniscrews after failure of the primary insertion for maxillary anchorage: Maxillary buccal area vs midpalatal suture [19] Watanabe H, Deguchi T, Hasegawa M, Ito M, Kim S, area. *Am J Orthod Dentofacial Orthop*. 2018;153(2):54-60.
- [29] Van Hevele J, Nout E, Claeys T, Meyns J, Scheerlinck J, Politis C. Bone-anchored maxillary protraction to correct a class III skeletal Bone-anchored maxillary protraction to correct a class III skeletal relationship: A multicenter retrospective analysis of 218 patients. *J Craniomaxillofac Surg*. 2018;46(10):1800-6.
- [30] Haddad R, Saadeh M. Distance to alveolar crestal bone: a critical factor in the success of orthodontic mini-implants. *Prog Orthod*. 2019;20(1):19-4.