Published Online: 2023-06-30

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



Mathematical Study of Weibull – New Weighted Exponential Distribution

Adetunji K. Ilori¹, Folakemi M. Okafor², Oladimeji D. Matthew³, Dayo V. Kayode⁴

¹Statistics Programme, National Mathematical Center, Abuja, Nigeria

²Mathematics Programme, National Mathematical Center, Abuja, Nigeria

^{3,4}Department of Statistics, University of Abuja, FCT, Nigeria

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— A four Parameter Weibull - New Weighted Exponential distribution was proposed with the use of the Weibull Generalized family of distributions. The Weibull New Weighted Exponential distribution has the ability to model failure rates which are quite common in medical sciences and reliability/survival Studies. Mathematical properties of the proposed distribution such as Moments, reliability analysis and order Statistics were derived. The Method of Maximum Likelihood Estimation was used in estimating the distribution Parameters. Keywords— Generalization, maximum likelihood estimation, Azallini, order statistics, Weibull Generalized family of distributions.

Introduction

Numerous classical distributions have been extensively considered by researcher for analysing lifetime data set over the past decades. Different extensions of existing distributions have been considered for modelling data in several areas of real life data such as, actuarial, medical sciences, engineering and many more. However, in many of these applied areas, there is continuous need for extended forms of existing distributions. For that reason, several methods for generating new families of distributions have been studied. (Bourguignon, Silva and Cordeiro, 2014)

The Weibull, gamma and exponential distributions are common distributions that have been extensively used over years for modelling data in reliability and biological studies. Much attention has been given to these distributions in literature in modelling lifetime data. However,

Published Online: 2023-06-30

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



researchers continue to develop different extensions of the Weibull, gamma and exponential distribution that will suit different behaviour of real life data set.

The Weibull distribution is a very popular model and the distribution has been considered extensively over the past years for modelling data in reliability and engineering studies. It is generally adequate for modelling monotone hazard rates. Bourguignon et al., (2014) Bourguignon et al., (2014) gave expression for generating Weibull family of distribution as follows: Consider a positive continuous distribution G with density g and the Weibull CDF

$$F(\mathbf{x}) = 1 - e^{-\alpha x^{\beta}}$$
 (for x>0) with positives α and β .

Based on this density, by replacing x with $\frac{G(\mathbf{x})}{1-G(\mathbf{x})}$, the

CDF family is defined by

$$F(x,\alpha,\beta) = \int_{0}^{\frac{G(x)}{1-G(x)}} \alpha \beta t^{\beta-1} e^{-\alpha t^{\beta}} dt$$

$$F(x,\alpha,\beta) = 1 - \exp\left\{-\alpha \left[\frac{G(x)}{1 - G(x)}\right]^{\beta}\right\} \qquad x,\alpha,\beta > 0$$

Where $G(\mathbf{x})$ CDF of baseline distribution and the family PDF is given as

$$f(x,\alpha,\beta) = \alpha\beta g(x) \frac{\left[G(x)\right]^{\beta-1}}{\left[1 - G(x)\right]^{\beta+1}} \exp\left\{-\alpha \left[\frac{G(x)}{1 - G(x)}\right]^{\beta}\right\}$$

3

By using the exponential function in (3), we have

$$\exp\left\{-\alpha \left[\frac{G(\mathbf{x})}{1-G(\mathbf{x})}\right]^{\beta}\right\} = \sum_{k=0}^{\infty} \frac{(-1)^{k} \alpha^{k}}{k!} \left[\frac{G(\mathbf{x})}{1-G(\mathbf{x})}\right]^{k\beta}$$

.

E-ISSN: 2833-7298, P-ISSN: 2833-7301



Aljarrah, Famoye and Lee, (2015) and Cordeiro, Edwin and 9 The cdf can further be Ramires, (2015) gave expressions for other forms of 10 generalizing Weibull family of distribution. Another expression of Equation (1) was given by Nasiru, and Luguterah, (2015) as follows

$$F(x) = \int_{0}^{\frac{1}{1-G(x)}} f(x)dx$$

A 3-parameter probability distribution called Weibull 11 Exponential distribution was proposed by Oguntunde, Balogun and Amina, (2015) using the concept of the Weibull Generalized family of distribution.

Materials And Methods

Let consider distribution proposed by Oguntunde et al (2016) called new weighted exponential distribution as the baseline distribution on the weibull family of distribution as expressed in Equation (2) and (3) The pdf and cdf of the new weighted exponential distribution by Oguntunde et al (2016) are defined as

$$g(x) = (1 + \lambda)\alpha e^{-(1+\lambda)\alpha x}$$

$$G(x) = 1 - e^{-(1+\lambda)\alpha x}$$

To derive the cdf of the Weibull new weighted exponential distribution, the cdf of the new weighted exponential distribution as given in Equation (8) is inserted into Equation (2). The resulting expression is given as:

$$F(x) = 1 - e \left[-\theta \left[\frac{1 - e^{-(1+\lambda)\alpha x}}{e^{-(1+\lambda)\alpha x}} \right]^{\beta} \right]$$

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



The cdf can further be simplify as:

$$F(x) = 1 - e^{-\theta \left[e^{(1+\lambda)\alpha x} - 1\right]^{\beta}}$$

Also, the corresponding pdf is obtained by inserting the pdf and cdf of the new weighted exponential distribution into

Equation (3). The resulting expression is given as:

$$f(x) = \alpha \beta (1+\lambda) \theta e^{-(1+\lambda)\alpha x} \left[\frac{(1-e^{-(1+\lambda)\alpha x})^{\beta-1}}{(e^{-(1+\lambda)\alpha x})^{\beta+1}} \right] e^{\left[-\theta \left[\frac{1-e^{-(1+\lambda)\alpha x}}{e^{-(1+\lambda)\alpha x}}\right]^{\beta}\right]}$$

The pdf can f_u rthe f_v be simplify f_v as:

$$f(x) = \alpha \beta \theta (1+\lambda) (1 - e^{-(1+\lambda)\alpha x})^{\beta-1} e^{(1+\lambda)\alpha\beta x - \theta [e^{(1+\lambda)\alpha x} - 1]^{\beta}}$$
12




Figure 1: cdf of Weibull new weighted exponential distribution

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



where,
$$a = \alpha$$
, $b = \beta$, $c = \theta$, $d = \lambda$




Figure 2: pdf of Weibull new weighted exponential x, distribution

$$\lambda = d\theta = c\beta = b\alpha = where, a$$

Statistical Properties Of Weibull New Weighted Exponential Distribution Reliability Analysis:

The Survivor function indicates the probability that the event of interest has not yet occurred by time t; thus, if T denotes probability of)t(denotes time until failure, S surviving beyond t.

The expression for Survival (or reliability) function is given as:

$$(S)x(F-1=)x(S)$$

Therefore, the survival function for the WNWE distribution is

$$S(x) = e^{-\theta [e^{(1+\lambda)\alpha x}-1]^{\beta}}$$
; $x,\alpha,\beta,\lambda,\theta > 0$

Hazard Function Of Wnwedistribution

Hazard function is a conditional probability that the event of interest (device failure) will occur during the time t and dt under the condition that the device is safe until time t.



hazard function is defined as the ratio of probability density given as:)x(to the survival function S)x(function f)

$$h(x) = \frac{f(x)}{S(x)}$$

Therefore, the hazard function for the WNWE distribution is:

$$h(x) = \frac{1}{\alpha\beta\theta (1+\lambda)(1-e^{-(1+\lambda)\alpha x})^{\beta-1}e^{(1+\lambda)\alpha\beta x-\theta [e^{(1+\lambda)\alpha x}]^{\beta}}}$$

$$x,\alpha,\beta,\lambda,\theta > 0 \qquad 14$$

Reversed Hazard Function Of Wnwe Distribution

The reverse hazard function is defined as the ratio of to the distribution)x(probability density function f of a random variable, given by <math>)x(function F)

$$r(x) = \frac{f(x)}{F(x)}$$

Therefore, the reversed hazard function for the WNWE distribution is:

$$r(x) = \frac{\alpha\beta\theta (1+\lambda) (1-e^{-(1+\lambda)\alpha x})^{\beta-1} e^{(1+\lambda)\alpha\beta x-\theta [e^{(1+\lambda)\alpha x}-1]^{\beta}}}{1-e^{-\theta [e^{(1+\lambda)\alpha x}-1]^{\beta}}}$$

$$; x,\alpha,\beta,\lambda,\theta > 0 15$$

Published Online: 2023-06-30

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



Moments

Some properties of the Weibull Generalized family of distributions can be directly obtained from those of the exponentiated Generalized family of distributions (Bourguignon et al., 2014). Therefore, the rth moment of X can be expressed as:

$$E(X^r) = \sum_{j,k} W_{j,k} E(Z_{jk})$$

Then, Weibull exponential distribution, Zj,k is the exponentiated exponential distribution with power $1-j+)+(k\beta parameter$

For a random sample X1.) $\beta,\lambda,\alpha,...$, Xn the probability density Equation (12), the Likelihood function $L(x \mid function \ of the ith order statistic from a distribution function <math>F(x)$ and probability density function f(x) is defined as

$$f_{i,n}(x) = \frac{n!}{(i-1)!(n-i)!} f(x) F(x)^{i-1} [1 - F(x)]^{n-i}$$

Inserting the pdf and cdf of the Weibull new weighted exponential distribution into the expression of probability density function of the ith order statistics. The resulting expression is given as

$$f(x) = \frac{n!}{(1+\lambda)(\alpha\beta x - \beta\beta x)(\alpha x - \beta\beta} tn - \lceil \alpha\beta \xi (1+\lambda)(1 - e^{-(1+\lambda)(\alpha x)})^{\beta - 1} \epsilon$$

$$(i-1)!(n-i)! \sum_{j=1}^{n} (1 - e^{-\beta \xi^{\beta \lambda}(\alpha x - \beta\beta})^{\beta - 1} \left[e^{-\beta \xi^{\beta \lambda}(\alpha x - \beta\beta})^{\beta - 1} \right]^{n-\beta}$$

Published Online: 2023-06-30

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



of Weibull)1(The pdf of the Minimum order statistic X gives the maximum likelihood estimates of the θ and model parameters. Also, the solution can be obtained directly by using R software when data sets are available. new weighted exponential distribution is obtained as

$$\begin{split} f_1\left(x\right) &= n \bigg[\alpha\beta\theta\left(1+\lambda\right)\left(1-e^{-(1+\lambda)\alpha x}\right)^{\beta-1}e^{(1+\lambda)\alpha\beta x-\theta\left[e^{(1+\lambda)\alpha x}-1\right]^{\beta}}\bigg] \\ &\times \bigg[e^{-\theta\left[e^{(1+\lambda)\alpha x}-1\right]^{\beta}}\bigg]^{n-1} \end{split}$$

Parameter Estimation

The Estimation of the Weibull new weighted exponential distribution is obtained using the Method of Maximum Likelihood Estimation (MLE) as follows:

Let x ,x2 ,...,xn be a random sample of size "n" from Weibull new weighted exponential distribution defined in

Equation (12), the Likelihood function L(x.) β , λ , α /I s given by

$$\begin{split} L(x/\alpha,\lambda,\beta,\theta) &= \prod_{i=1}^n f(\mathbf{x}_i/\alpha,\lambda) \\ \text{let } I &= \log L(x/\alpha,\lambda,\beta,\theta) \\ I &= n \log \alpha + n \log \beta + n \log \theta + n \log (1+\lambda) + (\beta-1) \sum_{i=1}^r \log [1-e^{-(1+\lambda)\alpha_{\lambda_i}}] \\ &+ \sum_{i=1}^n \{(1+\lambda)\beta\alpha x_i - \theta[e^{(1+\lambda)\alpha_{\lambda_i}} - 1]^\beta\} \end{split}$$

The solution of the non-linear system of equations obtained β , λ , α by differentiating Equation (21) with respect to 18 of Weibull)1(The pdf of the Minimum order statistic X gives the maximum likelihood estimates of the θ and model parameters. Also, the solution can be obtained directly by using R software when data sets are available

Published Online: 2023-06-30

International Journal of Multidisciplinary Research and Explorer (IJMRE)

E-ISSN: 2833-7298, P-ISSN: 2833-7301



Conclusion

A four-parameter Weibull new weighted exponential distribution has been successfully derived and defined. The proposed distribution can be suitable for modelling lifetime 19 data. The shape of model is unimodal. properties of proposed distribution such as survival function, hazard function, moment and order statistics are discussed. The method of maximum likelihood was considered for parameter estimation. The Weibull new weighted exponential distribution can be used as an alternative for a life testing model.

References

- [1] Aljarrah, M.A, Famoye, F. and Lee, C. 2015. A new weibull pareto distribution. Communications in statistics Theory and Methods, 44:19, 4077-4095
- [2] Bourguignon, M., R.B. Silva and G.M. Cordeiro, 2014. The weibull-G family of probability distributions. J. Data Sci., 12: 53-68.
- [3] Cordeiro, G.M, Ortega E. and Ramires T.G., 2015. A new generalized weibull family of distributions: Mathematical properties and applications. Journal of statistical distributions and applications
- [4] Nasiru, S. and A. Luguterah, 2015. The new weibull-pareto distribution. Pak. J. Stat. Operat. Res., 11: 103-114
- [5] Oguntunde PE, Balogun O.S and Amina B.S., 2015. The weibull Exponential distribution: Its properties and Applications. Journal of Applied Fire Science, 15(11): 1305-1311
- [6] Oguntunde, PE, Balogun O.S. and Owoloko A.E., 2016. On a new weighted exponential distribution: Theory and Applications. Asian journal of applied sciences, 9(1): 1-12